

# **Need to Know: A Fact Sheet Series on Juvenile Suicide**

## **Juvenile Detention and Secure Care Staff**

Prepared by the  
Youth in Contact With the Juvenile Justice System Task Force  
of the National Action Alliance for Suicide Prevention

September 2013





## Need to Know: A Fact Sheet Series on Juvenile Suicide Juvenile Detention and Secure Care Staff

### What You Need to Know about Juvenile Suicide

#### ✿ Youth suicide is a significant, yet preventable, public health problem.

- Suicide is the second leading cause of death among youth ages 10–18.<sup>1</sup>
- About one in thirteen high-school students attempted suicide in the past year.<sup>2</sup>
- Nearly 88,000 youth ages 10–18 were treated in emergency rooms for self-harm injuries in 2011.<sup>3,4</sup>
- Males are more likely to die by suicide<sup>1</sup> and females are more likely to attempt suicide.<sup>4</sup>
- Certain populations (e.g., American Indian/Alaskan Native and sexual minority youth) have increased rates of suicide.<sup>5,6</sup>

#### About this Fact Sheet

*Need to Know: A Fact Sheet Series on Juvenile Suicide* was developed by the Youth in Contact with the Juvenile Justice System Task Force of the National Action Alliance for Suicide Prevention. The task force's Public Awareness and Education Workgroup developed this series to raise awareness among individuals who work with youth involved with the juvenile justice system.

#### ✿ Youth involved with the juvenile justice system have *increased* risk of suicide.

- Suicide is the leading cause of death for youth in confinement.<sup>7</sup>
- Youth in residential facilities have nearly three times the suicide rate of peers in the general population.<sup>8</sup>
- Risk factors for suicide are often more prevalent among youth in the juvenile justice system.<sup>9</sup>
- Studies report that over half of juveniles had current suicidal ideation<sup>10</sup> and one-third had a history of suicidal behavior.<sup>11</sup>

#### ✿ Factors that may *increase* the risk of suicide among all youth include:<sup>11,12,13,14</sup>

- Mental illness and/or substance use disorder
- History of suicide attempts, self-harm behavior, and/or death by suicide in the family
- Social isolation, relationship problems, or separation from family
- Impulsive, aggressive, or reckless behavior
- History of bullying or being bullied
- Access to lethal means
- History of trauma or child maltreatment

#### ✿ Factors that may *increase* suicide risk among youth in the justice system include:<sup>15,16, 17,18</sup>

- History of mental illness and/or substance use disorder
- Involvement in special education
- Legal/disciplinary problems
- Prior disciplinary action
- Prior offenses
- Referral to juvenile court
- Being placed on room confinement



## **Need to Know:** **A Fact Sheet Series on Juvenile Suicide** **Juvenile Detention and Secure Care Staff**

---

- ✿ **Signs that *immediate* help for suicide risk is needed include:**<sup>10,11,12,13</sup>
  - Perceived crisis (e.g., transition within the juvenile justice system)
  - Unusual or sudden changes in personality, behavior, or mood
  - Talking about wanting to die or kill oneself
  - Withdrawal from friends, family, or usual activities
  - Expressions of hopelessness or feeling trapped
  - Actively securing access to lethal means
- ✿ **Protective factors that may *decrease* suicide risk among youth in the justice system include:**<sup>18,19,20</sup>
  - Easy access to effective mental health and substance abuse treatment services
  - Problem-solving and conflict-resolution skills
  - Cultural or religious beliefs that discourage suicide
  - Connectedness and support from family and community
  - A positive school experience
  - Lack of access to lethal means
  - Suicide-resistant housing
  - Collaborative communication between juvenile justice and mental health systems

### **What You Can Do to Prevent Juvenile Suicide**

- ✿ **Demonstrate your belief that suicide *can be* prevented.**<sup>15</sup>
- ✿ **Learn the protective factors, risk factors, and warning signs related to suicide.**<sup>15</sup>
- ✿ **Implement and evaluate comprehensive suicide prevention policies, programs, and practices that address risk and protective factors on multiple levels.**
- ✿ **Take any written, spoken, or other communication of suicide seriously. If you think someone is at risk:**
  - Do not be afraid to ask if someone is considering suicide. This *will not* cause suicide.
  - Be direct. Ask:
    - Are you thinking about killing yourself?
    - Are you considering taking your own life?
    - Do you ever feel like things would be better if you were dead?
  - Listen and do not judge anyone who you think might be thinking of suicide.
  - Avoid acting shocked if a youth says he or she is considering suicide.
  - Do not be sworn to secrecy or make promises that you won't tell anyone.
  - Communicate any suspicion that a youth is thinking about suicide to a mental health professional or supervisor *immediately*.
  - Stay with the youth. Do *not* leave a suicidal youth alone while you go get help.



**Need to Know:**  
**A Fact Sheet Series on Juvenile Suicide**  
**Juvenile Detention and Secure Care Staff**

---

**What Detention and Secure Care Facilities Can Do to Prevent Juvenile Suicide**

✿ **Develop, implement, and maintain a comprehensive written suicide prevention program that includes the following eight critical components:<sup>13</sup>**

- Routine suicide prevention training for all staff
- Standardized intake screening for suicide risk using a valid and reliable tool for all youth, with suicide risk assessment by a qualified mental health professional administered as necessary
- Protocols that provide shared information about suicide risk
  - Among the arresting/transporting officer, family members, and facility staff
  - Between facility staff members
  - Between facility staff and youth
- Varying levels of supervision<sup>21</sup>
  - Close observation for youth with suicidal ideation or behavior
  - Constant observation for youth who are talking about or displaying suicidal behavior

\*\*Closed-circuit television does **not** substitute for observation.
- Safe physical environment
- Emergency response protocol in the case of suicides or suicide attempts
- Notification system for suicides or suicide attempts through the chain of command
- Critical incident stress debriefing protocol (for all staff and youth) and a death review

✿ **Access additional resources for more information.**

- Centers for Disease Control and Prevention
  - <http://www.cdc.gov/ViolencePrevention/suicide>
- National Center on Institutions and Alternatives, Inc.
  - <http://www.ncianet.org/services/suicide-prevention-in-custody>
- National Suicide Prevention Lifeline
  - <http://www.suicidepreventionlifeline.org>; (800) 273-TALK (8255)
- Office of Juvenile Justice and Delinquency Prevention
  - <http://www.ojjdp.gov>
- Substance Abuse and Mental Health Services Administration
  - <http://www.samhsa.gov/prevention/suicide.aspx>
- Suicide Prevention Resource Center
  - <http://www.sprc.org>



## Need to Know: A Fact Sheet Series on Juvenile Suicide Juvenile Detention and Secure Care Staff

### About This Fact Sheet

*Need to Know: A Fact Sheet Series on Juvenile Suicide* was developed by the Youth in Contact with the Juvenile Justice System Task Force (<http://actionallianceforsuicideprevention.org/task-force/juvenilejustice>) of the National Action Alliance for Suicide Prevention (<http://actionallianceforsuicideprevention.org>) (Action Alliance). The task force's Public Awareness and Education Workgroup developed this series to raise awareness among individuals who work with youth involved with the juvenile justice system, as well as provide practical information on what:

- **you need to know** about juvenile suicide
- **you can do** to prevent juvenile suicide
- **systems can do** to prevent juvenile suicide

This fact sheet is tailored to juvenile detention and secure care staff; companion pieces are tailored to juvenile probation staff and to juvenile court judges and staff.

The Public Awareness and Education Workgroup included the following members and staff:

- Deborah Stone, ScD, MSW, MPH (*workgroup lead*) – Behavioral Scientist, National Center for Injury Prevention and Control, Centers for Disease Control and Prevention
- Christy Lentz, MSW – Senior Policy Associate, National Association of State Mental Health Program Directors
- Roy Praschil – Director of Operations, National Association of State Mental Health Program Directors
- Steffie Rapp, LCSW-C – Program Manager, Office of Juvenile Justice and Delinquency Prevention, U.S. Department of Justice
- Kathleen Skowrya – Associate Director, National Center for Mental Health and Juvenile Justice, Policy Research Associates

*Suggested citation:* National Action Alliance for Suicide Prevention: Youth in Contact with the Juvenile Justice System Task Force. (2013.) *Need to know: A fact sheet series on juvenile suicide – juvenile detention and secure care staff*. Washington, DC: Author.

### Action Alliance

Envisioning a nation free from the tragic experience of suicide, the Action Alliance was launched in 2010 by U.S. Department of Health and Human Services Secretary Kathleen Sebelius and former U.S. Department of Defense Secretary Robert Gates. This public-private partnership advances the *National Strategy for Suicide Prevention* (NSSP) by championing suicide prevention as a national priority, catalyzing efforts to implement high-priority objectives of the NSSP, and cultivating the resources needed to sustain progress.

The Action Alliance established the Youth in Contact with the Juvenile Justice System Task Force to focus attention on the needs of youth in the juvenile justice system. The task force was co-led by:

- Melodee Hanes, JD – Acting Administrator, Office of Juvenile Justice and Delinquency Prevention, Office of Justice Programs, U.S. Department of Justice
- Joseph J. Coccozza, PhD – Director, National Center for Mental Health and Juvenile Justice, Policy Research Associates

The task force comprised four workgroups: Public Awareness and Education; Suicide Research; Suicide Prevention Programming and Training; and Mental Health and Juvenile Justice Systems Collaboration.



## Need to Know: A Fact Sheet Series on Juvenile Suicide Juvenile Detention and Secure Care Staff

### References

- <sup>1</sup> Centers for Disease Control and Prevention, National Center for Injury Prevention and Control. Web-based Injury Statistics Query and Reporting System (WISQARS). *Fatal injury data*. Accessed November 2012. Retrieved from <http://www.cdc.gov/injury/wisqars/fatal.html>
- <sup>2</sup> Centers for Disease Control and Prevention. *1991–2011 High school youth risk behavior survey data*. Accessed August 2012. Retrieved from <http://apps.nccd.cdc.gov/youthonline>
- <sup>3</sup> The term “self-harm” includes suicide attempts and non-suicidal self-injuries.
- <sup>4</sup> Centers for Disease Control and Prevention, National Center for Injury Prevention and Control. Web-based Injury Statistics Query and Reporting System (WISQARS). *Nonfatal injury data*. Accessed October 2012. Retrieved from <http://www.cdc.gov/injury/wisqars/nonfatal.html>
- <sup>5</sup> Centers for Disease Control and Prevention, National Center for Injury Prevention and Control. Web-based Injury Statistics Query and Reporting System (WISQARS). *Fatal injury data*. Accessed August 2012. Retrieved from <http://www.cdc.gov/injury/wisqars/fatal.html>
- <sup>6</sup> Suicide Prevention Resource Center. (2008). *Suicide risk and prevention for lesbian, gay, bisexual and transgender youth*. Newton, MA: Education Development Center, Inc.
- <sup>7</sup> Bureau of Justice Statistics. *Deaths in custody statistical tables: state juvenile correctional facility deaths, 2002–2005*. Retrieved from <http://bjs.ojp.usdoj.gov/content/dcrp/tables/juvtab1.cfm>
- <sup>8</sup> Gallagher, C. A. & Dobrin, A. (2006). Deaths in juvenile justice residential facilities. *Journal of Adolescent Health* 38: 662–668.
- <sup>9</sup> National Action Alliance for Suicide Prevention: Youth in Contact with the Juvenile Justice Task Force – Suicide Research Workgroup. (2013). *Suicidal ideation and behavior among youth in the juvenile justice system: A review of the literature*. Washington, DC: Author.
- <sup>10</sup> Esposito, C., & Clum, G. (2002). Social support and problem-solving as moderators of the relationship between childhood abuse and suicidality: Applications to a delinquent population. *Journal of Traumatic Stress*, 15(2), 137–146.
- <sup>11</sup> Hayes, L. (2009). Characteristics of juvenile suicide in confinement. *Juvenile Justice Bulletin*. Washington, DC: U.S. Department of Justice, Office of Juvenile Justice and Delinquency Prevention. Retrieved from <https://www.ncjrs.gov/pdffiles1/ojdp/214434.pdf>
- <sup>12</sup> Gould, M.S., Greenberg, T., Velting, D.M. & Shaffer, D. (2003). Youth suicide risk and preventive interventions: A review of the past 10 years. *Journal of the American Academy of Child and Adolescent Psychiatry* 42: 386–405.
- <sup>13</sup> Beautrais, A.L. (2000). Risk factors for suicide and attempted suicide among young people. *The Australian and New Zealand Journal of Psychiatry* 34(3): 420–436.
- <sup>14</sup> Brent, D.A., Baugher, M., Bridge, J., Chen, T. & Chiappetta, L. (1999). Age- and sex-related risk factors for adolescent suicide. *Journal of the American Academy of Child & Adolescent Psychiatry* 38(12): 1497–1505.
- <sup>15</sup> Gray, D., Moskos, M. (2005). Utah youth suicide study phase VI: Treatment for those at highest risk for suicide in Utah-Juvenile Offenders. Webinar for the Suicide Prevention Resource Center. Retrieved from <http://www.sprc.org/training-institute/r2p-webinars/utah-youth-suicide-study-phase-vi-treatment-those-highest-risk-suici>
- <sup>16</sup> Hayes, L. (2004). *Juvenile suicide in confinement: A national survey*. Washington, DC: Office of Juvenile Justice and Delinquency Prevention. Retrieved from <https://www.ncjrs.gov/pdffiles1/ojdp/213691.pdf>
- <sup>17</sup> Gray, D. et al. (2002). Utah youth suicide study phase I: Government agency contact before death. *Journal of the American Academy of Child and Adolescent Psychiatry*, 41(4), 427–434.
- <sup>18</sup> Kaminski, J.W., Puddy, R.W., Hall, D.M., Cashman, S.Y., Crosby, A.E. & Ortega, L.A. (2010). The relative influence of different domains of social connectedness on self-directed violence in adolescence. *Journal of Youth and Adolescence* 39(5): 460–473.
- <sup>19</sup> U.S. Public Health Service. (1999). *The Surgeon General’s call to action to prevent suicide*. Washington, DC: U.S. Department of Health and Human Services.
- <sup>20</sup> Kidd, S., Henrich, C.C., Brookmeyer, K.A., Davidson, L., King, R.A. & Shahar, G. (2006). The social context of adolescent suicide attempts: interactive effects of parent, peer, and school social relations. *Suicide & Life-threatening Behavior* 36(4): 386–395.
- <sup>21</sup> National Action Alliance for Suicide Prevention: Youth in Contact with the Juvenile Justice System Task Force. (2013). *Guide to developing and revising suicide prevention protocols for youth in contact with the juvenile justice system*. Washington, DC: Author.

The National Action Alliance for Suicide Prevention is the public-private partnership advancing the *National Strategy for Suicide Prevention* (NSSP) (<http://actionallianceforsuicideprevention.org/NSSP>) by championing suicide prevention as a national priority, catalyzing efforts to implement high-priority objectives of the NSSP, and cultivating the resources needed to sustain progress. The Action Alliance envisions a nation free from the tragic experience of suicide. For electronic copies of this paper or for additional information about the Action Alliance and its task forces, please visit <http://www.actionallianceforsuicideprevention.org>.

